

Review of glucose control, GI and type 2 diabetes with a focus on fruit juice

1. Glycemic response

Glycemic response refers to the effect that foods and drinks have on blood glucose after consumption. After eating a meal, carbohydrates (excluding fibre) are absorbed from the intestine into the bloodstream, causing a temporary increase in blood glucose concentration – this is called a glucose excursion. In response, the hormone insulin is released and blood glucose concentration returns to fasting levels, or may fall slightly below.

There is clear scientific evidence that persistent elevation of blood glucose is linked with conditions such as type 2 diabetes, cardiovascular disease and obesity. A systematic review and metaanalysis confirmed that diets with a low glycemic impact (i.e. resulting in only moderate elevations of blood glucose) have a role in reducing the risk of such chronic conditions¹.

There are several markers for assessing glycemic status, all of which are used to diagnose diabetes². Normal and abnormal values are shown in Table 1.

- Fasting blood glucose levels
- Oral Glucose Tolerance Test (OGTT), which measures how quickly a 75 g dose of glucose is cleared from the blood in a fasting individual
- Concentrations of glycated proteins e.g. HbA1c, which reflect the impact of long-term elevations in blood glucose.

Table 1: Glycemic control markers criteria³

Marker	Normal	Diabetes
Fasting glucose	70–99 mg/dl (3.9–5.5 mmol/l)	≥ 126 mg/dl (7.0 mmol/l)
OGTT	< 140 mg/dl (7.8 mmol/l)	≥ 200 mg/dl (11.1 mmol/l)
HbA1c	< 6.5%	≥ 6.5 %

The glycemic effect of foods also depends on the sensitivity of tissues to insulin, a hormone secreted by the pancreas. When released, this stimulates insulin-sensitive tissues, such as muscle and adipose tissue, to take up glucose from the blood. As blood glucose returns to normal, insulin release is slowed to normal resting levels.

In cases of ‘insulin resistance’, muscle and adipose tissue do not respond adequately to insulin production. This results in a reduced rate of glucose uptake from blood with the consequence that blood glucose levels remain elevated. More insulin is then released leading to chronically high plasma insulin levels. In the long term, there is a risk that the pancreas may become exhausted, resulting in falling insulin levels and a need for insulin administration.

The homeostatic model assessment (HOMA) is often used to assess the risk of insulin resistance. This marker shows the dynamic between baseline (fasting) blood glucose and the responsive hormone insulin; the healthy range is 0.5–1.4.

2. Glycemic index (GI)

Carbohydrates provide a significant energy source in our diets. The glycemic response depends on the type of carbohydrate and its molecular composition. In addition, GI is influenced by the food matrix in which the carbohydrates are present. The matrix, in turn, is influenced by food processing and the presence of dietary fibre, protein and fat. Carbohydrate-containing foods that result in a slow increase in blood glucose may be called low glycemic foods.

Pure glucose is rapidly taken up by the bloodstream


and is given a standard glycemic value of 100. By comparing carbohydrate foods to this standard of 100, it is possible to give these a relative index score. In this respect, GI is defined as the area under the glucose response curve after consumption of 50 g carbohydrate from a test food divided by the area under the curve after consumption of 50 g glucose (sometimes white bread is taken as standard, instead of glucose)⁴.

Generally, there are three categories of foods based on their GI values⁵:

- high GI foods (≥ 70)
- intermediate GI foods (56–69)
- low GI foods (≤ 55)

Consuming low GI foods, instead of high GI foods, has a positive effect on keeping post-meal glucose excursions and related insulin requirements relatively low (Fig. 1). Fully and readily digestible carbohydrates, such as glucose, maltodextrin, white bread and cooked potato starch, produce a rapid increase in blood glucose and insulin, followed by an equally rapid fall. If high GI foods are eaten in significant amounts, this has implications for the overall glycemic impact of the meal. Consuming high glycemic meals frequently in conditions of overweight and inactivity will drive the development of insulin resistance and type 2 diabetes.


Figure 1: Blood glucose response in healthy adults⁶



Legend: After a high glycemic meal there is a rapid increase in blood glucose. This triggers a significant insulin release which stimulates tissues to take up glucose from blood. As a result, the blood glucose level will fall rapidly, sometimes even below the pre-meal level (hypoglycemic effect). In contrast, consumption of a low glycemic meal will produce a moderate increase in blood glucose and insulin, and a slow return to the pre-meal blood glucose level.

Consumption of fructose, for example found in fruits, table sugar, honey and fruit juices, only minimally increases blood glucose, explaining why fructose-containing foods tend to have a relatively low glycemic response. This is also the reason why 100% fruit juices have a low GI, contrary to popular belief and media claims that they produce a significant rise in blood glucose (Fig 2.)⁷.

Figure 2: Examples of plasma glucose curves (mmol/l) after different food challenges



3. Glycemic load (GL)

The glycemic response to a food depends not just on GI but also on the total carbohydrate ingested. On this basis GL was created; defined by how much each portion of carbohydrate raises blood glucose levels. GL is classified as: low (≤ 10), intermediate (11–19) and high (≥ 20). Fruits and 100% fruit juices have a low GI and intermediate GL (Table 2).

Table 2: GI/GL of typical products⁸

Food item	GI/100 g	GL/portion
White bread	75+2	11/30 g
Whole wheat bread	74+2	7/30 g
Cornflakes	81+6	21/30 g
White rice, boiled	73+4	28/150 g
Apple, raw	36+2	6/120 g
Orange, raw	43+3	4/120 g
Orange juice, medium	50+2	11/250 ml
Orange juice, small	50+2	7/150 ml
Banana, raw	51+3	11/120 g
Potato, boiled	78+4	21/150 g
Sugar-sweetened drinks	63-68	16-23/250 ml

A systematic review and meta-analysis¹ of data from 45 studies reported that diets comprised of low glycemic foods reduced fasting blood glucose and HbA1c, particularly in people with poor fasting blood glucose control. Adding non-digestible carbohydrates enhanced these effects. Lower GL was more important than GI for lowering triglyceride levels.

4. 100% fruit juice, glycemic control and type 2 diabetes

In one trial, 36 overweight subjects with raised plasma cholesterol were enrolled in a randomised, single-blinded, placebo-controlled clinical study⁹ to investigate the metabolic effects of daily 100% orange juice consumption. During the 12-week intervention, participants received either 250 ml of orange juice or a control orange drink matched for energy and sugars. The results revealed that 100% orange juice had no adverse effects on insulin sensitivity (HOMA-IR), blood lipid profile or body weight. The authors concluded that: "Daily consumption of 250 ml of orange juice for three months did not result in an increase in

dietary sugars intake in a cohort of overweight men with elevated total cholesterol concentration, and, despite media concern, an increase in body weight or decreased insulin sensitivity did not occur over the intervention".

A meta-analysis¹⁰ of 12 randomised controlled trials involving over 400 participants who were obese or had risk factors for diabetes or cardiovascular disease examined the effect of sugar-containing beverages on fasting glucose and insulin levels. In half of these studies, the intake of 100% fruit juice was 400 ml per day or more. The overall results showed that consumption of 100% fruit juices had no significant effect on fasting blood glucose or insulin levels.

Another meta-analysis¹¹ of four cohorts of adults found that consumption of fruit drinks with added sugars was significantly associated with an increased risk of type 2 diabetes (RR = 1.28; $p = 0.02$) while consumption of 100% fruit juices (i.e. no added sugars) had no effect (RR = 1.03, $p = 0.62$).

The most recent systematic review and meta-analysis¹², based on 18 randomised and controlled studies, examined the effects of 100% fruit juice on glucose-insulin homeostasis. Compared with controls, 100% fruit juices had no significant effect on fasting blood glucose (mean difference: -0.13 mmol/l; 95% CI $-0.28, 0.01$; $p=0.07$), fasting blood insulin (-0.24 mmol/l; 95% CI $-3.54, 3.05$; $p=0.89$), HOMA-IR (-0.22 ; 95% CI $-0.50, 0.06$; $p=0.13$) or HbA1c (-0.001% ; 95% CI $-0.38, 0.38$; $p=0.28$). The authors concluded that their meta-analysis suggested: "a neutral effect of 100% fruit juice on glycemic control" and, consequently, "consumption of 100% fruit juice is not associated with increased risk of diabetes".

Conclusion

Overall, the available evidence shows clearly that 100% fruit juices have no negative effects on glucose-insulin homeostasis and are not a causal factor in the development of type 2 diabetes. This probably reflects the fact that 100% fruit juices have a low GI and contain significant amounts of bioactive components, such as flavanones, that have been shown to exhibit lipid-lowering, insulin-sensitising, antihypertensive and anti-inflammatory properties¹³. This may also explain the observation that a high intake of 100% orange juice and sugar-sweetened beverage, matched for energy and sugars, differently affects metabolic risk in healthy subjects¹⁴.

References

1. Livesey G, et al. (2008) Glycemic response and health – a systematic review and meta-analysis: relations between dietary glycemic properties and health outcomes. *Am J Clin Nutr* 87: 258S–268S.
2. American Diabetes Association (2014) Standards of medical care in diabetes. *Diabetes Care* 37: S14–80.
3. American Diabetes Association (2015) Standards of medical care in diabetes. *Diabetes Care* 38: 58–67.
4. Brouns F et al. (2005) Glycaemic index methodology. *Nutr Res Rev* 18: 145–171.
5. Glycemic Index Foundation www.gisymbol.com/about-glycemic-index.
6. Harvard Health (2012) Choosing good carbs with the glycemic index www.health.harvard.edu/staying-healthy/choosing-good-carbs-with-the-glycemic-index.
7. Brand-Miller J et al. (2009) Glycaemic index, postprandial glycemia, and the shape of the curve in healthy subjects: analysis of a database of more than 1000 foods. *Am J Clin Nutr* 89: 97–105.
8. Atkinson RD et al. (2008) International Tables of Glycemic Index and Glycemic Load Values. *Diabetes Care* 31: 2281–2283.
9. Simpson EJ et al. (2016) Orange juice consumption and its effect on blood lipid profile and indices of the metabolic syndrome; a randomised, controlled trial in an at-risk population. *Food Funct* 7: 1884–91.
10. Wang B et al. (2014) Effect of fruit juice on glucose control and insulin sensitivity in adults: a meta- analysis of 12 randomized controlled trials. *PLoS ONE* 9: e95323.
11. Xi B et al. (2014) Intake of fruit juice and incidence of type 2 diabetes: a systematic review and meta- analysis. *PLoS ONE* 9: e93471.
12. Murphy MM et al. (2017) 100% Fruit juice and measures of glucose control and insulin sensitivity: a systematic review and meta-analysis of randomised controlled trials. *J Nutr Sci* 6: e59 (15 pages).
13. Mulvihill EE et al. (2016) Citrus Flavonoids as Regulators of Lipoprotein Metabolism and Atherosclerosis. *Annu Rev Nutr* 36: 275–99.
14. Büsing F et al. (2018) High intake of orange juice and cola differently affects metabolic risk in healthy subjects, *Clin Nutr*; epub ahead of print: <https://doi.org/10.1016/j.clnu.2018.02.028>.